

Ultraviolet dominance of power corrections in QCD?

V.M. Braun

NORDITA, Blegdamsvej 17, DK-2100 Copenhagen Ø, Denmark

Presented at 5th International Conference on Physics Beyond the Standard Model,
Balholm, Norway, 29 Apr - 4 May 1997

Abstract

I explain current theoretical ideas on higher-twist (hadronization) corrections to physical observables in QCD.

The recent years witnessed an outburst of theoretical activity devoted to nonperturbative effects in hard processes in Quantum Chromodynamics. For a generic physical observable R one considers the perturbative expansion complemented by nonperturbative corrections which are expected to be suppressed by a power of the hard scale Q

$$R(Q, \underline{x}) = R_0(\underline{x}) \left[1 + r_1(\underline{x}) \frac{\alpha_s(Q)}{\pi} + r_2(\underline{x}) \left(\frac{\alpha_s(Q)}{\pi} \right)^2 + \dots \right] + \frac{A_R^{(p)}(\underline{x})}{Q^p} \quad (1)$$

Here $\underline{x}$ is a set of dimensionless variables (typically momentum fractions), R_0 is the tree-level perturbative result, r_i give the radiative corrections and $A_R^{(p)}$ is a (nonperturbative) constant of dimension GeV^p determining the leading power correction (with the smallest power of Q). Interest in power suppressed contributions is due to our desire to make QCD predictions (in the perturbative domain) as quantitative as possible, which is imperative for a precise determination of the parameters of the Standard Model and the ongoing searches of “new physics”. The experimental accuracy has reached the level at which power like effects are observable and can be studied. This calls for a broad and systematic study of the theory and phenomenology of nonperturbative effects in hard processes in QCD, for which little was done until very recently.

The main obstacle for the phenomenology has been an almost complete absence of theoretical predictions, with the Monte Carlo hadronization models providing the only

approach to the data. This situation has changed and we have a “minimal model” of higher-twist effects which is consistent with QCD, sufficiently general, simple enough and introduces a minimum number of parameters. This model can be directly confronted with the data; both agreement and disagreement would be meaningful and interesting. This is still not a theory, but an “educated guess” to shoot at. The objective of my talk today is to emphasize this recent development which is based on the examination of the structure of infrared contributions in perturbation theory in somewhat more detail than usual in the derivations of celebrated factorization theorems.

To explain the basic idea, I will consider power corrections to the structure functions of deep inelastic scattering. This example provides a benchmark for the theoretical understanding since the Operator Product Expansion (OPE) allows to express power corrections to moments of the structure functions in terms of contributions of higher twist operators. Schematically, one can write

$$\int dx x^{N-1} F_2(x, Q^2) = C(N, \mu) P(N, \mu) + \frac{1}{Q^2} \sum_k C_k^{(4)}(N, \mu) P_k^{(4)}(N, \mu) + O\left(\frac{1}{Q^4}\right) \quad (2)$$

where $P(N)$ are the moments of quark and gluon parton distributions, $C(N)$ are the corresponding coefficient functions, while $C_k^{(4)}$ and $P_k^{(4)}$ are the coefficient functions and matrix elements of the relevant twist 4 operators, respectively. The summation goes over the complete set of such operators which are rather numerous [1].

From the OPE one learns that the nonperturbative effects are suppressed as $1/Q^2$ and are given in terms of universal objects (in fact, moments of multiparton correlation functions [2]) with calculable coefficients in perturbation theory. Thus, power corrections to different deep inelastic processes are related; in principle one can sacrifice a few measurements to extract higher-twist distributions and then predict the corrections to other processes. In practice, however, the multiparton distributions are too complicated; thus predictive power of the OPE is largely lost. As a result, despite being available for 15 years, the results of [1, 2] on higher twist parton distributions have rarely been used in phenomenological analysis.

I am going to formulate a certain approximation and start from the observation that the separation of leading twist from higher twists is not unique. Imagine that the factorization in transverse momenta that is implicit in (2) were implemented by a rigid cutoff μ such that only coefficients from transverse momenta $k_t > \mu$ were included in the coefficient function $C(N, \mu)$. The logarithmic cutoff dependence would cancel between the coefficient function and the parton distribution, as usual. In addition, one would find, however, a *power-like* cutoff dependence starting from μ^2/Q^2

$$C(N, \mu) = c_0(N) + \frac{\alpha_s}{\pi} \left[c_1(N, \log \mu) + h_1(N, \log \mu) \frac{\mu^2}{Q^2} + \dots \right] + \dots \quad (3)$$

which is cancelled by higher-twist contributions. This cancellation is possible since matrix elements of twist four operators are quadratically divergent with the scale μ serving as an *ultraviolet* cutoff and it means that the coefficient in front of this quadratic divergence

must match the IR cutoff dependence of the coefficient function

$$\sum_k C_k^{(4)}(N, \mu) P_k^{(4)}(N, \mu) / P(N, \log \mu) = \mu^2 \frac{\alpha_s}{\pi} h_1(N, \log \mu) + A^{(4)}(N, \log \mu) \quad (4)$$

Thus the leading twist contribution as a whole depends on the prescription used to implement the scale separation, just as the separation of the leading twist coefficient function and parton distribution does. At first sight, such prescription dependence seems to be avoided in the $\overline{\text{MS}}$ scheme, because power-like dependence on the factorization scale μ does not exist. The problem reappears, however, because the coefficient function now has a factorially divergent series expansion in α_s (referred to as infrared renormalon divergence). Summing the series again requires a prescription and the prescription-dependence is power suppressed precisely as the cutoff dependence above. In both cases, the ultraviolet renormalization of higher-twist operators must be performed consistently with the definition of the leading-twist coefficient function. The sum of leading-twist and higher-twist contributions is then unique.

Now comes the central point. As it is well known, the logarithmic scale dependence does not in fact cancel exactly in finite-order perturbative calculations, and provides a conventional estimate for the accuracy of the calculation. This implies a tacit assumption that the scale independent constant terms in the next (uncalculated) order of the perturbation theory are of the same order of magnitude as scale dependent logarithmic terms which can easily be reconstructed from the requirement that they must exactly cancel the calculated scale dependence in low orders. Thus, although the scale dependent terms are ultimately unphysical and must cancel, their magnitude indicates the size of uncalculated higher-order corrections. Very similarly, the power like scale dependence of perturbation theory in (3) can be used to estimate the size of higher-twist corrections under the assumption that the “true” matrix elements $A^{(4)}$ in (4) are roughly proportional to their quadratically divergent pieces

$$A^{(4)}(N, \log \mu) \simeq \mu_0^2 h_1(N, \log \mu) \quad (5)$$

where μ_0 is a certain scale of order Λ_{QCD} . Note that μ_0 is assumed to be (approximately) independent of N ; Thus, the N dependence of higher-twist corrections is tied up with (calculable) N dependence of IR contributions in perturbation theory. I will refer to this assumption as ultraviolet dominance of power corrections [3].

Since the coefficient function C in (2) does not depend on the target (by definition), so does the ratio of the higher twist correction to the leading twist parton distribution in (4). The ultraviolet dominance approximation should be viewed, therefore, as a “minimal model” of supposedly large generic target-independent higher twist corrections, providing an “irreducible background” for “true” effects of quark-gluon nonperturbative correlations. Note that from the phenomenological point of view the higher-twist correction can only be defined as a sum of “true” nonperturbative effects and perturbative corrections beyond available order.

Calculation of the characteristic functions $h_1(N)$ in leading-order radiative corrections is relatively easy and can be done using different techniques [4, 5, 6, 7]. The results exist for all flavor non-singlet deep inelastic structure functions and for the polarized structure

function g_1 [7, 8, 9]. A preliminary comparison to the existing data was done in [10] (see also [7]), with encouraging results. Similar calculations for flavor-singlet contributions are in progress [11].

Extension of these results beyond leading order faces difficulties, both technical and conceptual, which I do not have time to discuss in this talk.

Since the approach is purely perturbative it can be generalized to a variety of other processes where the OPE does not exist. The inclusive cross section of hadron production in e^+e^- annihilation [12, 3] provides an instructive example. Tracing IR contributions $\sim \mu^2/Q^2$ in the leading twist coefficient function we get an ansatz for the power corrections to the longitudinal and total cross sections [3]

$$\begin{aligned} \frac{d\sigma_L^{\text{power}}}{dx}(x, Q^2) = & \frac{\mu_0^2}{Q^2} \int_x^1 \frac{dz}{z} \left\{ c_{q,L} \left[\delta(1-z) + \frac{2}{z} \right] D_q(x/z, \mu) \right. \\ & \left. + c_{g,L} \frac{1-z}{z^3} D_g(x/z, \mu) \right\}, \end{aligned} \quad (6)$$

$$\begin{aligned} \frac{d\sigma_{L+T}^{\text{power}}}{dx}(x, Q^2) = & \frac{\mu_0^2}{Q^2} \int_x^1 \frac{dz}{z} \left\{ c_{q,L+T} \left[-\frac{2}{[1-z]_+} + 1 + \frac{1}{2} \delta'(1-z) \right] D_q(x/z, \mu) \right. \\ & \left. + c_{g,L+T} \frac{1-z}{z^3} D_g(x/z, \mu) \right\}, \end{aligned} \quad (7)$$

where x is the energy fraction carried by the detected hadron and D_i denotes the leading-twist fragmentation function for parton $i = q, g$. The parametrization depends on four constants c_q, c_g which have to be fitted from the data or related to some nonperturbative parameters (see below). With the overall scale factor $\mu_0 = 1$ GeV these constants are expected to be of order unity. One can check [13] that the expressions in (6), (7) correspond to ultraviolet contributions to multiparton fragmentation functions introduced in [14].

The most interesting potential application of these methods is to hadronic event shape observables. The major result in this case is that nonperturbative corrections to generic observables are large, of the order $1/Q$ [15, 16, 17, 18], and are due to soft gluon emission. Since the soft emission exponentiates, so does the leading nonperturbative correction [17]. This leads to specific predictions for event shape distributions [19] which appear to be in agreement with the data.

Most of the specific predictions have been obtained so far using the concept of the universal effective coupling [16, 7]. In this language nonperturbative corrections to physical cross sections are due to nonperturbative contributions to the effective coupling and are parametrized by its moments. In practice, application of this technique amounts to choosing the same IR parameter μ_0^2 in the relation (5) for all physical observables (but different for corrections of different power). This implies universality of nonperturbative scales for Euclidian and Minkowskian observables which would be very interesting to test experimentally.

To summarize, there are ongoing efforts to understand the structure of nonperturbative corrections to hard processes in QCD by using their relation to IR effects in perturbation theory. In case that the operator structure of the higher twists is known, these corrections can be identified as being due to ultraviolet regions in the relevant matrix elements, which explains the title of my talk. The resulting expressions should be viewed as theoretically motivated parametrizations of higher twist corrections by a few phenomenological parameters.

A striking feature of these parametrizations is that they are target-independent, if weighted by leading twist distributions as explained above. This approximation tacitly assumes absence of any enhancements of higher twists due to strong quark-gluon correlations in hadrons and is supported by estimates of higher twist corrections to deep inelastic sum rules (see a discussion in [20]) which appear to be of the same order of magnitude as renormalon ambiguities in perturbation theory. It is also supported by the recent analysis in [21] which suggests that phenomenological extractions of higher twist corrections essentially parametrize higher order perturbative rather than nonperturbative effects. Consider, however, deep-inelastic scattering at small x from a heavy nucleus. The main higher twist corrections are due in this case to Glauber-type rescattering effects which have nothing to do with the leading twist perturbation theory. The ultraviolet dominance approximation would miss the rescattering effects (and thus the A dependence) completely. The reason is precisely that the large number of nucleons provides a new parameter which governs the hierarchy of higher twist effects.

The moral is that the significance of target-dependent vs. target-independent higher twist effects has to be studied experimentally, which provides a very particular task for the phenomenology. On the other hand, the theory in its present form is still far from being complete. One problem is that IR contributions to gluon fragmentation are not calculated directly, but are related to quark contributions of particular type (chains of bubbles) or to the dependence on a certain naive IR regulator like the gluon mass. This identification is not exact for insufficiently inclusive observables, which causes ambiguities in predictions, see [3] for a detailed discussion. This is related to the observation that universality of IR contributions for event shape observables is spoiled by emission of soft gluons at large angles which is process-dependent [22]. A more general problem is that the strong coupling in (4) stands at a low scale μ_0 which indicates that all orders of perturbation theory are formally important (unless one can prove the opposite by calculation of the anomalous dimension of the relevant higher twist operators). A discussion of these (and other) difficulties as well as further theoretical ideas on power corrections and related issues can be found in recent reviews [23].

References

- [1] Shuryak E.V., and Vainshtein A.I., *Nucl. Phys.* **B199**, 451 (1982);
Jaffe R.L., and Soldate M., *Phys. Rev.* **D26**, 49 (1982).

- [2] Ellis R.K., Furmanski W., and Petronzio R., *Nucl. Phys.* **B212**, 29 (1983);
Jaffe R.L., *Nucl. Phys.* **B229**, 205 (1983);
Balitsky I.I., and Braun V.M., *Nucl. Phys.* **B311**, 541 (1988/89).
- [3] Beneke M., Braun V.M., and Magnea L., *Nucl. Phys. Proc. Suppl.* **54A**, 183 (1997) (hep-ph/9609266); *Nucl. Phys.* **B497**, 297 (1997).
- [4] Beneke M., *Nucl. Phys.* **B405**, 424 (1993).
- [5] Beneke M., Braun V.M., and Zakharov V.I., *Phys. Rev. Lett.* **73**, 3058 (1994).
- [6] Beneke M., and Braun V.M., *Phys. Lett.* **B348**, 513 (1995);
Ball P., Beneke M., and Braun V.M., *Nucl. Phys.* **B452**, 563 (1995).
- [7] Dokshitzer Yu.L., G. Marchesini G., and Webber B.R., *Nucl. Phys.* **B469**, 93 (1996).
- [8] Stein E. et al., *Phys. Lett.* **B376**, 177 (1996); Meyer-Hermann M. et al., *Phys. Lett.* **B383**, 463 (1996); *Erratum-ibid.* **B393**, 487 (1997).
- [9] Dasgupta M., and Webber B.R., *Phys. Lett.* **B382**, 273 (1996).
- [10] Maul M. et al., *Phys. Lett.* **B401**, 100 (1997).
- [11] Mankiewicz L., *private communication*.
- [12] Dasgupta M., and B.R. Webber B.R., *Nucl. Phys.* **B484**, 247 (1997).
- [13] Beneke M., Braun V.M., and Magnea L., *work in progress*.
- [14] Balitsky I.I., and Braun V.M., *Nucl. Phys.* **B361**, 93 (1991).
- [15] Webber B.R., *Phys. Lett.* **B339**, 148 (1994).
- [16] Dokshitzer Yu.L., and Webber B.R., *Phys. Lett.* **B352**, 451 (1995).
- [17] Korchemsky G.P., and Sterman G., *Nucl. Phys.* **B437**, 415 (1995); Contribution to 30th Rencontres de Moriond: QCD and High Energy Hadronic Interactions, March 1995 (hep-ph/9505391).
- [18] Akhouri R., and Zakharov V.I., *Phys. Lett.* **B357**, 646 (1995); *Nucl. Phys.* **B465**, 295 (1996).
- [19] Dokshitzer Yu.L., and Webber B.R., CAVENDISH-HEP-97-2, April 1997 (hep-ph/9704298).
- [20] Braun V.M., Contribution to 30th Rencontres de Moriond: QCD and High Energy Hadronic Interactions, March 1995 (hep-ph/9505317).
- [21] Kataev A.L. et al., Contribution to 32nd Rencontres de Moriond: QCD and High-Energy Hadronic Interactions, March 1997 (hep-ph/9706534).
- [22] Nason P., and Seymour M.H., *Nucl. Phys.* **B454**, 291 (1995).
- [23] Akhouri R., and Zakharov V.I., *these proceedings* (hep-ph/9705318); Beneke, M., CERN-TH-97-134 (hep-ph/9706457); Korchemsky G.P., Oderda G., and Sterman G., ITP-SB-97-41 (hep-ph/9708346).